

Journal of Thermal Analysis and Calorimetry

Kinetic analysis of the thermal decomposition of liquid ammonium nitrate based on thermal analysis and detailed reaction simulations

--Manuscript Draft--

Manuscript Number:	JTAC-D-17-01589R1	
Full Title:	Kinetic analysis of the thermal decomposition of liquid ammonium nitrate based on thermal analysis and detailed reaction simulations	
Article Type:	S.I. : CEEC-TAC4	
Corresponding Author:	Yu-ichiro Izato, Ph. D Yokohama National University Yokohama, JAPAN	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Yokohama National University	
Corresponding Author's Secondary Institution:		
First Author:	Yu-ichiro Izato, Ph. D	
First Author Secondary Information:		
Order of Authors:	Yu-ichiro Izato, Ph. D Atsumi Miyake, Ph.D.	
Order of Authors Secondary Information:		
Funding Information:	Japan Society for the Promotion of Science (17H00844)	Professor Atsumi Miyake
Abstract:	<p>A detailed reaction mechanism for the liquid-phase decomposition of ammonium nitrate (AN) was modeled based on ab initio calculations, and this model was employed to simulate heat flow curves associated with the decomposition of AN at various heating rates (1, 2, 4, 5 and 8 K min⁻¹) over the temperature range of 443-623 K. A kinetic analysis using the model-free Friedman method determined the kinetic triplet for AN decomposition. The predicted activation energy for the exothermic decomposition had a range of approximately 162-168 kJ mol⁻¹ with α values between 0.1 and 0.5. The activation-energy value is in good agreement with the experimental one of 170 kJ mol⁻¹. The Friedman kinetic model was used to predict the time to maximum rate under adiabatic conditions (TMRad) at various initial temperatures, and these values were compared to the TMRad values obtained directly using the detailed reaction mechanism. Accurate predictions for TMRad were obtained at initial temperatures below 623 K. The difference between the TMRad values obtained from the thermal analysis and detailed reaction models at 823 K was larger than one order of magnitude. It was also found that the dominant decomposition mechanism changes from ionic to radical with increasing temperature, and this explains the difference in the TMRad values at 823 K.</p>	
Response to Reviewers:	Please see attachment.	

Kinetic analysis of the thermal decomposition of liquid ammonium nitrate based on thermal analysis and detailed reaction simulations

Yu-ichiro Izato

Graduate school of Environment and Information Sciences, Yokohama National University, 79-7 Tokiwadai, Hodogaya-ku, Yokohama, Kanagawa 240-8501, Japan

Phone: +81-45-339-3981

E-mail: izato-yuichiro-hk@ynu.jp

Atsumi Miyake

Institute of Advanced Sciences, Yokohama National University, 79-5, Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

Abstract A detailed reaction mechanism for the liquid-phase decomposition of ammonium nitrate (AN) was modeled based on *ab initio* calculations, and this model was employed to simulate heat flow curves associated with the decomposition of AN at various heating rates (1, 2, 4, 5 and 8 K min⁻¹) over the temperature range of 443–623 K. A kinetic analysis using the model-free Friedman method determined the kinetic triplet for AN decomposition. The predicted activation energy for the exothermic decomposition had a range of approximately 162–168 kJ mol⁻¹ with α values between 0.1 and 0.5. The activation-energy value is in good agreement with the experimental one of 170 kJ mol⁻¹. The Friedman kinetic model was used to predict the time to maximum rate under adiabatic conditions (TMR_{ad}) at various initial temperatures, and these values were compared to the TMR_{ad} values obtained directly using the detailed reaction mechanism. Accurate predictions for TMR_{ad} were obtained at initial temperatures below 623 K. The difference between the TMR_{ad} values obtained from the thermal analysis and detailed reaction models at 823 K was larger than one order of magnitude. It was also found that the dominant decomposition mechanism changes from ionic to radical with increasing temperature, and this explains the difference in the TMR_{ad} values at 823 K.

Keywords ammonium nitrate, thermal decomposition, kinetics analysis, time to maximum rate (TMR_{ad})

Introduction

Ammonium nitrate (AN) is widely used as a fertilizer ingredient because it is relatively inexpensive. Unfortunately, tragic accidental explosions involving AN have occurred in the past [1-4], including an incident at a West Fertilizer Company storage facility in Texas in 2013 that killed 14 and injured 260 [4]. Following this accident and others, the use and storage of AN have been strictly regulated. This background emphasizes the need to improve our ability to prevent accidental AN explosions. AN is also used as an oxidizer in industrial explosives because it releases almost 100% gaseous products upon reaction and has a positive oxygen balance (+20.0 g g⁻¹). In addition, AN has been considered as an alternative to ammonium perchlorate as a solid rocket propellant oxidizer [5]. However, the poor combustion properties of AN, including low ignitability and sluggish burning rates under low pressure conditions, have prevented the application of AN-based propellants to date. As a result, there have been many studies aimed at improving the combustion characteristics of AN in combination with various catalysts or fuels [5-12]. Despite this prior work, the combustion properties of AN-based propellants still require improvement.

Based on the above, it is evident that both the chemical and physical stability of AN must be increased to prevent unintended explosions during storage while, in contrast, good ignitability and high burning rates are needed to develop AN-containing propellants. To allow the safe development and use of AN-containing devices, it is important to understand both the reaction mechanisms and the kinetics controlling the combustion and decomposition properties. There have been many studies of the AN decomposition mechanism and a number of reasonable schemes have been proposed and summarized in the literature [5]. The decomposition mechanism of AN has been shown to have two potential pathways, based on ionic or radical reactions. The former path involves various ions and proceeds relatively slowly at low temperatures (< 563 K) [1, 13], while the latter involves active radicals, proceeds more rapidly, and is predominant at high temperatures (> 563 K) [13]. It is known that AN melts at 442 K and begins to decompose after it melts. The first step of the low temperature decomposition process involves the dissociation of NH₄NO₃ (NH₄⁺ and NO₃⁻) into NH₃ and HNO₃. The next step involves the self-decomposition of HNO₃ into NO₂⁺, NO₃⁻ and H₂O, followed by the oxidation of NH₃ by the NO₂⁺ [1]. Park and Lin [14] and Skarlis et al. [15] have also proposed another decomposition path that proceeds via NH₃OH⁺, while Manelis et al. [8] have suggested that HNO₃ oxidizes NH₄⁺ directly. In systems with a large excess of HNO₃, the predominant reaction changes to oxidation by N₂O₅. Our previous work [16] analyzed the ionic decomposition of AN in the liquid phase, using computations based on quantum mechanics to confirm the identity of products observed in past experimental studies. During these calculations, the CBS-QB3//wB97XD/6-311++G(d,p) method [17, 18] was employed. It was found that one of the most reasonable reaction pathways is HNO₃ + NH₄⁺ → NH₃NO₂⁺ + H₂O followed by NH₃NO₂⁺ + NO₃⁻ → NH₂NO₂ + HNO₃. The other path is HNO₃ + HNO₃ → N₂O₅ + H₂O followed by N₂O₅ + NH₃ → NH₂NO₂ + H₂O, after which the nitramide (NH₂NO₂) decomposes to N₂O and H₂O. The thermal decomposition of AN in the liquid phase can therefore be summarized as NH₄⁺ + NO₃⁻ (AN) → N₂O + 2H₂O. Although the proposed details of the AN decomposition reaction differ among researchers, the overall condensed-phase decomposition may be expressed as NH₄NO₃ → N₂O + 2H₂O. The gases resulting from the ionic reactions are primarily N₂O and H₂O, with N₂ as a minor product [19, 20]. As the temperature increases, the ionic decomposition of AN is thought to be overtaken by high temperature radical decomposition. and Brower et al. [13] suggested a mechanism for the radical reaction of AN at elevated temperatures (> 563 K). In this mechanism, AN initially dissociates into HNO₃ and NH₃, followed by homolytic cleavage of the HO-N bond in HNO₃. Following this cleavage, a high-speed radical chain reaction develops and yields large amounts of gaseous N₂, NO and H₂O. Since the activation energy for the homolysis of HNO₃ is very high (approximately 190 kJ mol⁻¹), this represents the rate controlling step and radical reactions do not play an important role at low temperatures (< 563 K) [13].

Thermal analysis is a powerful tool for the investigation of the decomposition kinetics of various materials. Kinetic analysis can have either a theoretical or a practical purpose. The theoretical purpose of kinetic analysis is typically interpretation of experimentally-determined kinetic triplets, while the most common practical purpose is the prediction of process rates and material lifetimes. The goal of kinetic analysis is to parameterize the process rate in terms of variables such as temperature, extent of conversion and, in some cases, pressure. Parameterization is accomplished by evaluating the parameters of equations that describe the effects of variables on the process rate. To date, there have been many studies of AN decomposition kinetics. Kinetic analysis based on thermal analysis is a useful means of examining the kinetics of AN

1 decomposition, and much research regarding AN has been performed under various conditions,
2 with the results summarized in reference [21]. The AN thermal decomposition kinetics depend on
3 several factors, including temperature, pressure, isothermal or non-isothermal conditions, extent of
4 reaction, catalysis and reactive species. The activation energies reported by various researchers
5 range of 30–200 kJ mol⁻¹, depending on the specific sample analyzed and the experimental
6 methods. Willis et al. [19] investigated the kinetics of decomposition of liquid AN (AN (L) → N₂O
7 (G) + 2H₂O (G)) in the temperature range 498 to 548 K using a flow reactor, and also reported the
8 activation energy of liquid phase decomposition is 170 kJ mol⁻¹.

9 In the present work, we examined both the theoretical and practical purposes of kinetic analysis
10 based on detailed reaction simulations. The specific goals were: (i) to simulate the AN
11 decomposition thermal behavior based on a detailed reaction model, (ii) to analyze the kinetics
12 using the model-free Friedman method so as to compare the kinetics obtained in reference [19],
13 and (iii) to simulate thermal behavior at various temperatures based on the kinetic analysis and
14 detailed reaction model. Kinetic analysis can also simulate the thermal behaviors of materials
15 under extreme conditions that cannot readily be achieved during thermal analysis. The resulting
16 predictions are reliable only when sound kinetic model involving adequate kinetic parameters (the
17 activation energy, the pre-exponential factor, and the reaction are used. This work verifies the
18 kinetics based on the model-free Friedman method, and compares the predictions from the kinetic
19 analysis and the detailed reaction simulation.

21 Computational

22 Detailed reaction model

23 The detailed reaction model consists of two parts: a collection of elementary reactions with their
24 rate coefficients and thermodynamic data. The present work employed the YNU-L 1.0 model,
25 consisting of various kinetic parameters (a total of 54 reactions) and the thermodynamic data for
26 29 species.

27 The rate coefficient, k_{TST} , of the generic reaction $A + B \rightarrow$ products can be calculated on the
28 basis of traditional transition state theory (TST) using the well-known formula $k_{\text{TST}} =$
29 $\frac{k_{\text{B}}T}{h} \frac{Q_{\text{TS}}}{\prod Q_{\text{react}}} \exp\left(\frac{-\Delta E_0}{RT}\right)$, where k_{B} is the Boltzmann constant, T is the temperature, h is the Planck
30 constant, Q_i is the partition function of the reactant and transition state (TS), ΔE_0 is the energy
31 barrier to activation, and R is the universal gas constant. The TST provides rate of an elementary
32 reaction, if we obtain Q_i and ΔE_0 of TS and reactants. Both of Q_i and ΔE_0 can be computed by
33 quantum chemical calculation. In our previous work [16], liquid-phase reactions were investigated
34 based on quantum chemical calculation at the CBS-QB3 [17] //ωB97XD [18] /6-
35 311++G(d,p)/SCRF = (solvent = water) level of theory and the associated potential energy
36 diagrams, reactants, and TS structures were identified and investigated. The TS is a surface in
37 configuration space that divides reactants from products and passes through the saddle point of the
38 potential-energy surface. In this study, we calculated k_{TST} of reactions identified in our previous
39 study [16, 22]. Variational TS theory (VTST) was applied to the analysis of dissociation reactions
40 without barriers. The VTST is distinguished by varying the definition of the TS to minimize the
41 one-way rate coefficient. These calculations were aided by the GPOP software package developed
42 by Miyoshi [23]. Radical recombinations and proton transfers with no barriers were modeled as
43 diffusion-limited reactions with rate coefficients set at 10⁹ cm³ mol⁻¹ s⁻¹. The important reactions
44 associated with the decomposition of liquid-phase AN and the associated kinetic parameters for the
45 modified Arrhenius equation are provided in Table 1 (for the ionic pathway) and Table 2 (for the
46 radical pathway).

47 Thermal correction, entropy (S_{liq}) and heat capacity (C_p) values were calculated from the Q_i
48 using statistical machinery, employing the GPOP software package [23]. The heats of formation
49 for gas-phase molecules ($\Delta_f H_{\text{gas}}^\circ$) were calculated by the traditional atomization method (ARM-1)
50 [24] combined with the G4 [25]/SCRF = (solvent = water) level of theory using the Gaussian 09
51 program package [26]. Solvent effects were included by applying the self-consistent reaction field
52 (SCRF) and polarizable continuum model (PCM) options within the program when investigating
53 the liquid species in molten AN. Unfortunately, the solvent effect of molten AN is not known. The
54 dielectric constant for solid AN has been reported to be approximately 40 [-] at 383 K [27], and it
55 is known that this value tends to increase along with temperature. Thus, molten AN could be
56 considered as a highly polar solvent, so we employed the water solvation effect as a substitute for
57 molten AN. It is noted that the water solvation effect should be replaced with a more adequate

solvation effect in future work. In the interim, we believe that the use of water solvation is a practical means of obtaining some insights regarding the liquid-phase decomposition of AN.

The standard heat of formation for a compound in solution is obtained from the gas-phase heat of formation and the enthalpy of solvation at 298.15 K, as in the following two equations.

$$\Delta_f H_{liq}^\circ = \Delta_f H_{gas}^\circ + \Delta_{solv} H^\circ \quad (1)$$

$$\Delta_{solv} H^\circ = H_{liq,calc} - H_{gas,calc} \quad (2)$$

Here, $\Delta_f H_{liq}^\circ$ is the heats of formation for liquid phase compounds, $\Delta_f H_{gas}^\circ$ is the heats of formation for gas phase compounds, $\Delta_{solv} H^\circ$ is the solvation enthalpy, and $H_{liq,calc}$ and $H_{gas,calc}$ are the heats of formation at 298.15 K calculated directly using G4 methods. Table 3 lists the thermodynamic data for liquid-phase compounds obtained at the G4 level of theory.

Table 1. Important reactions and associated rate coefficients employed during the kinetic modeling of ionic decomposition.

No.	Reaction	$k = A \cdot T^n \exp(-\Delta E_a/RT)$		
		A^a	n	ΔE_a^b
1	$\text{AN} \rightleftharpoons \text{NH}_3 + \text{HNO}_3$	3.03×10^{16}	-0.95	41.1
2	$\text{AN} \rightleftharpoons \text{NH}_4^+ + \text{NO}_3^-$	6.50×10^{20}	-2.01	65.1
3	$\text{NH}_3 + \text{HNO}_3 \rightleftharpoons \text{NH}_2\text{NO}_2 + \text{H}_2\text{O}$	1.22×10^1	3.11	168.7
4	$\text{NH}_4^+ + \text{HNO}_3 \rightleftharpoons \text{NH}_3\text{NO}_2^+ + \text{H}_2\text{O}$	4.50×10^2	3.66	123.1
5	$\text{AN} + \text{HNO}_3 \rightleftharpoons \text{NH}_2\text{NO}_2 + \text{HNO}_3 + \text{H}_2\text{O}$	1.16×10^1	3.58	123.1
6	$\text{HNO}_3 + \text{HNO}_3 \rightleftharpoons \text{N}_2\text{O}_5 + \text{H}_2\text{O}$	2.19×10^2	3.20	89.9
7	$\text{NH}_3 + \text{N}_2\text{O}_5 \rightleftharpoons \text{NH}_2\text{NO}_2 + \text{HNO}_3$	1.72×10^2	2.98	22.2
8	$\text{NH}_4^+ + \text{N}_2\text{O}_5 \rightleftharpoons \text{NH}_3\text{NO}_2^+ + \text{HNO}_3$	5.20×10^3	3.26	88.5
9	$\text{NH}_3\text{NO}_2^+ + \text{NO}_3^- \rightleftharpoons \text{NH}_2\text{NO}_2 + \text{HNO}_3$	1.00×10^9	0.0	0.0
10	$\text{NH}_3\text{NO}_2^+ + \text{NH}_3 \rightleftharpoons \text{NH}_2\text{NO}_2 + \text{NH}_4^+$	1.00×10^9	0.0	0.0
11	$\text{HNO}_3 + \text{HONO} \rightleftharpoons \text{t-ONONO}_2 + \text{H}_2\text{O}$	9.45×10^{-1}	3.60	5.3
12	$\text{t-ONONO}_2 \rightleftharpoons \text{NO}_2 + \text{NO}_2$	2.95×10^{12}	0.17	33.1
13	$\text{t-ONONO}_2 + \text{NH}_3 \rightleftharpoons \text{NH}_2\text{NO} + \text{HNO}_3$	1.00×10^9	0.0	0.0
14	$\text{t-ONONO}_2 + \text{NH}_4^+ \rightleftharpoons \text{NH}_3\text{NO}^+ + \text{HNO}_3$	1.44×10^2	3.21	60.6
15	$\text{NH}_3\text{NO}^+ + \text{NO}_3^- \rightleftharpoons \text{NH}_2\text{NO} + \text{HNO}_3$	1.00×10^9	0.0	0.0
16	$\text{NH}_3\text{NO}^+ + \text{NH}_3 \rightleftharpoons \text{NH}_2\text{NO} + \text{NH}_4^+$	1.00×10^9	0.0	0.0
17	$\text{NH}_2\text{NO}_2 \rightleftharpoons \text{NHNO}_2\text{H}$	2.10×10^{-1}	3.20	128.7
18	$\text{NH}_2\text{NO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{NHNO}_2\text{H} + \text{H}_2\text{O}$	1.47	3.09	53.6
19	$\text{NHNO}_2\text{H} \rightleftharpoons \text{N}_2\text{O} + \text{H}_2\text{O}$	8.77×10^9	1.23	134.9
20	$\text{NHNO}_2\text{H} + \text{H}_2\text{O} \rightleftharpoons \text{N}_2\text{O} + \text{H}_2\text{O} + \text{H}_2\text{O}$	5.46	3.58	99.8
21	$\text{NHNO}_2\text{H} + \text{NO}_3^- \rightleftharpoons \text{N}_2\text{O} + \text{H}_2\text{O} + \text{NO}_3^-$	7.52×10^2	3.41	35.6
22	$\text{NHNO}_2\text{H} + \text{NH}_3 \rightleftharpoons \text{N}_2\text{O} + \text{H}_2\text{O} + \text{NH}_3$	2.03×10^{-1}	3.34	-2.6
23	$\text{H}_3\text{O}^+ + \text{OH}^- \rightleftharpoons \text{H}_2\text{O} + \text{H}_2\text{O}$	1.00×10^9	0.0	0.0
24	$\text{NH}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$	1.00×10^9	0.0	0.0
25	$\text{NH}_4^+ + \text{H}_2\text{O} \rightleftharpoons \text{NH}_3 + \text{H}_3\text{O}^+$	1.00×10^9	0.0	0.0
26	$\text{HNO}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NO}_3^- + \text{H}_3\text{O}^+$	1.00×10^9	0.0	0.0
27	$\text{NO}_3^- + \text{H}_2\text{O} \rightleftharpoons \text{HNO}_3 + \text{OH}^-$	1.00×10^9	0.0	0.0

^a Frequency factor A is given in units of cm^3 , mol, and s.

^b Activation energy ΔE_a is in units kJ mol^{-1} .

Table 2. Reactions and associated rate coefficients employed during the kinetic modeling of radical decomposition.

No.	Reaction	k		
		A^a	n	ΔE_a^b
28	$\text{HNO}_3 \rightleftharpoons \text{NO}_2 + \text{OH}$	4.63×10^{12}	1.40	181.6
29	$\text{NH}_3 + \text{OH} \rightleftharpoons \text{NH}_2 + \text{H}_2\text{O}$	9.09×10^2	3.10	10.3
30	$\text{NH}_3 + \text{NO}_2 \rightleftharpoons \text{NH}_2 + \text{HONO}$	5.46	3.58	99.8
31	$\text{NH}_3 + \text{NO}_2 \rightleftharpoons \text{NH}_2 + \text{HNO}_2$	1.92×10^1	3.54	125.1
32	$\text{NH}_2 + \text{NO}_2 \rightleftharpoons \text{NH}_2\text{NO}_2$	1.00×10^9	0.0	0.0
33	$\text{NH}_2 + \text{NO}_2 \rightleftharpoons \text{NH}_2\text{ONO}$	2.00×10^3	2.81	-15.0
34	$\text{NH}_2\text{ONO} \rightleftharpoons \text{NH}_2\text{O} + \text{NO}$	9.45×10^{12}	0.30	54.1
35	$\text{NH}_2 + \text{NO} \rightleftharpoons \text{NH}_2\text{NO}$	1.00×10^9	0.0	0.0
36	$\text{NH}_2\text{NO} \rightleftharpoons \text{NHNOH}$	2.15×10^{-1}	3.82	98.4
37	$\text{NHNOH} \rightleftharpoons \text{N}_2 + \text{H}_2\text{O}$	1.76×10^{11}	0.89	99.4
38	$\text{NHNOH} + \text{H}_2\text{O} \rightleftharpoons \text{N}_2 + \text{H}_2\text{O} + \text{H}_2\text{O}$	4.60	3.96	54.6
39	$\text{NHNOH} + \text{NH}_3 \rightleftharpoons \text{N}_2 + \text{H}_2\text{O} + \text{NH}_3$	6.55	3.65	-1.7
40	$\text{NHNOH} + \text{NH}_3 \rightleftharpoons \text{N}_2 + \text{H}_2\text{O} + \text{NH}_3$	5.49	3.96	4.1
41	$\text{NH}_2\text{NO} + \text{H}_2\text{O} \rightleftharpoons \text{NHNOH} + \text{H}_2\text{O}$	6.02×10^{-3}	3.79	40.4
42	$\text{NH}_2\text{O} + \text{NO}_2 \rightleftharpoons \text{HNO} + \text{HONO}$	3.25×10^1	2.98	32.6
43	$\text{HNO}_2 + \text{NO}_2 \rightleftharpoons \text{HONO} + \text{H}_2\text{O}$	7.54×10^1	3.05	56.2
44	$\text{HNO}_2 + \text{HNO}_2 \rightleftharpoons \text{HONO} + \text{HONO}$	1.89×10^{-4}	4.42	21.8

^a Frequency factor is given in units of cm^3 , mol, and s.

^b Activation energy is in units kJ mol^{-1} .

Table 3. Calculated thermodynamic values for species associated with AN decomposition.

SPECIES	$\Delta_f H_{liq}^\circ$ [kJ/mol]	S_{liq}° [J/mol K]	Cp [J/K mol]						
			300 K	400	500	600	800	1000	1500
H ₂ O	-257.6	194.5	33.44	34.02	34.91	35.95	38.2	40.57	45.98
NO	88.1	205.2	29.16	29.44	30.00	30.74	32.26	33.49	35.30
NO ₂ ·	26.6	239.9	36.95	39.99	42.95	45.56	49.45	51.96	55.08
N ₂ O	74.6	219.3	37.97	41.94	45.08	47.66	51.57	54.28	58.04
N ₂ O ₅	-6.2	344.9	91.59	105.27	115.86	124.01	135.13	141.87	150.02
rONONO ₂	16.5	340.3	81.65	91.70	99.61	105.80	114.42	119.78	126.39
HNO ₃	-154.4	271.1	52.44	62.13	69.87	75.90	84.32	89.80	97.52
HONO	-92.1	253.2	44.34	50.33	55.17	59.02	64.59	68.42	74.25
HNO ₂	-68.8	238.1	38.26	43.51	48.84	53.59	61.00	66.19	73.64
HNO	95.3	220.6	33.68	34.95	36.86	38.99	43.04	46.31	51.42
NH ₃	-55.8	198.1	34.69	37.19	40.24	43.35	49.24	54.54	64.68
NH ₂ ·	173.6	194.7	33.53	34.25	35.30	36.50	39.15	41.87	47.53
AN	-281.9	336.7	89.51	103.68	116.73	128.14	146.15	159.23	179.22
NH ₂ O	37.4	225.4	34.59	37.22	40.42	43.57	49.08	53.59	61.51
NH ₂ NO ₂	-24.5	274.0	57.33	68.90	78.42	86.04	97.10	104.69	116.12
NHNO ₂ H	20.3	273.9	57.38	69.35	79.07	86.73	97.70	105.17	116.40
NH ₂ ONO	76.2	286.8	65.28	74.46	82.34	88.89	98.82	105.91	116.86
NH ₂ NO	55.2	258.9	49.89	57.24	63.75	69.19	77.45	83.43	92.99
NHNOH	62.5	254.6	46.90	55.15	62.28	68.11	76.82	83.02	92.80
NH(OH)NO	50.9	284.3	65.38	75.05	83.24	89.92	99.74	106.59	117.10
HNOHNO	96.6	263.8	58.06	68.60	77.85	85.53	96.98	104.94	116.73
N ₂	-2.4	191.5	29.11	29.20	29.47	29.93	31.16	32.37	34.48
OH·	24.4	178.2	29.1	29.10	29.12	29.19	29.58	30.26	32.30
H ₃ O ⁺	-107.2	198.8	35.41	38.05	41.00	43.92	49.39	54.35	64.17
NH ₄ ⁺	-53.0	191.5	34.63	38.14	43.01	48.19	57.96	66.41	81.74
NH ₃ NO ₂ ⁺	126.5	284.7	63.97	75.70	86.09	94.92	108.64	118.67	134.43
NH ₃ NO ⁺	163.2	278.7	61.69	69.28	75.84	81.53	90.95	98.47	111.46
NO ₃ ⁻	-184.0	245.5	44.43	52.29	58.81	63.85	70.54	74.43	78.95
OH ⁻	-96.5	172.3	29.10	29.10	29.13	29.21	29.62	30.34	32.40

Detailed reaction simulation

The YNU-L 1.0 mechanism was employed to simulate the heat flow curves for AN decomposition under non-isothermal conditions in an adiabatic reactor (at constant enthalpy and volume). These calculations were performed with the CHEMKIN-PRO software package

[28]. The initial density was set to 1.725 g cm⁻³, which is the density of pure liquid AN [5, 6], and the decomposition reactions were simulated at heating rates of 1, 2, 4, 5 and 8 K min⁻¹ (the heating rates typically used in thermal analysis) from 443 to 623 K. The resulting heat flow curves were investigated using the Friedman method as showed below.

The YNU-L 1.0 mechanism was also employed to predict the temperature rise in an adiabatic reactor (at constant enthalpy and volume) when applying initial temperatures of 423, 523, 623 and 823 K. The temperature rise data obtained in this manner were compared to those generated from kinetic analysis based on the Friedman method.

Kinetic analysis and thermal behavior prediction

The basic equation for kinetic analysis is:

$$\frac{d\alpha}{dt} = A(\alpha) \exp\left(-\frac{E(\alpha)}{RT(t)}\right) f(\alpha) \quad (3)$$

where α is the reaction progress, t is the time, $f(\alpha)$ is the reaction model, $A(\alpha)$ is the pre-exponential factor (note that both $f(\alpha)$ and $A(\alpha)$ are functions of the reaction progress), R is the gas constant, T is the temperature of the materials (T is constant in isothermal tests or a function of time in non-isothermal tests) and $E(\alpha)$ is the apparent activation energy.

The kinetic procedure employed in this study was based on the differential iso-conversion method of Friedman [29] and Ozawa [30]. The formula associated with the Friedman method (Eq. (4)) is obtained by the rearrangement of Eq. (3).

$$\ln\left(\frac{d\alpha}{dt}\right) = \ln A(\alpha)f(\alpha) - \frac{E(\alpha)}{RT(t)} \quad (4)$$

The values of α and $d\alpha/dt$ are determined from thermal analysis based on the following relationships.

$$\alpha = \frac{\int_{t_0}^t (S(t)-B(t))dt}{\int_{t_0}^{t_{end}} (S(t)-B(t))dt} \quad (5)$$

$$\frac{d\alpha}{dt} = \frac{S(t)-B(t)}{\int_{t_0}^{t_{end}} (S(t)-B(t))dt} \quad (6)$$

Here, $S(t)$ is a differential-type signal obtained from thermal analysis techniques such as differential scanning calorimetry (DSC) or differential thermogravimetric analysis (DTG) and $B(t)$ is the $S(t)$ baseline. For a given set of $d\alpha/dt$ values, a plot of $\ln(d\alpha/dt)$ values acquired at different heating rates as a function of $1/T$ can be fitted to a straight line, the slope of which gives the apparent activation energy. This method permits estimation of $E(\alpha)$ without knowing $f(\alpha)$ (model-free). The Friedman method is often used for the analysis of data obtained from scanning tests performed at a constant heating rate. However, this approach is based on differential kinetic law and therefore it can be applied to results from scanning or isothermal evaluations [29, 30]. In this study, the Friedman method was employed to elaborate the heat flow curves from detailed reaction simulations based on the YNU-L1.0 model. The AKTS thermokinetics software was used during kinetic analysis to provide a model-free Friedman kinetic model. This model simulated the adiabatic temperature rise at initial temperatures of 423, 523, 623 and 823 K, assuming a constant specific heat capacity of 1.5 K g⁻¹J⁻¹.

Results and discussions

Figure 1 shows the heat flow curves obtained from calculations for a heating rate of 5 K min⁻¹ using the YNU-L1.0 model as well as the experimental results acquired using a P-DSC apparatus [20, 31]. A heating furnace of the DSC can be pressurized up using any gases which is supplied from external compressed gas cylinders or gas supply lines. In our previous

1 studies [20, 31], Heating furnace was pressurized by helium gas at 1.1 MPa, and the gases in
2 furnace purged to external of it. The P-DSC results demonstrated that AN exhibits an
3 exothermic reaction above approximately 500 K with evolving N₂O, H₂O and minor N₂ gases
4 at pressures [20, 31]. The calculated sum of the moles of AN, NH₄⁺ and NH₃, indicating the
5 unreacted moles of AN, is plotted in Figure 2. This value begins to decrease at approximately
6 500 K, based on the criteria of a 1% reduction from the initial value. AN is known to undergo
7 exothermic decomposition between 473 and 503 K [5], and so this model successfully
8 predicts the decomposition onset temperature. Figure 1 also demonstrates that the moles of
9 the major products N₂O and H₂O increase beginning at this same onset temperature, and that
10 the final quantity of H₂O is twice that of N₂O. Omitting minor products, the reaction can be
11 summarized as AN → N₂O + 2H₂O. This new model therefore predicts the same evolved
12 gases as reported in previous studies [19, 20]. Thus, we concluded that the YNU-L 1.0 model
13 accurately predicted the decomposition behavior of AN, taking into account that the detailed
14 kinetics do not include any adjustable or empirical parameters. However, the initial calculated
15 heat flows were found to be higher than the experimental values, and the detailed model did
16 not predict the endothermic peak in experimental DSC curve following the exothermic peak
17 maximum. One reason for these deviations could be physical changes of the specimen,
18 including evaporation and gasification, since AN melts at 442 K and begins to gasify as soon
19 as it melts. The first step of the decomposition involves the dissociation of NH₄NO₃ into NH₃
20 and HNO₃, followed by the endothermic evaporation of NH₃ and HNO₃: AN(l) → NH₃(g) +
21 HNO₃(g) -174 kJ mol⁻¹. Other evolved gases will also evaporate from the molten AN. These
22 endothermic phase changes occur in the molten AN, offsetting the exothermic reaction. The
23 YNU-L1.0 model does not include such physical changes and so the experimental heat flow
24 data prior to the peak top will be lower than the calculated values.

25 Figure 3 presents the simulated heat flow curves obtained at heating rates of 1, 2, 4 and 8 K
26 min⁻¹. The average heat of reaction was 1940 ± 39 J g⁻¹, which is higher than the value of
27 1182 J g⁻¹ obtained from previous sealed DSC experiments [1]. This discrepancy is also
28 attributed to the neglect of physical changes in the detailed kinetic model. These heat flow
29 curves were elaborated by the Friedman method.

30 Figure 4 shows iso-conversion plots obtained from the heat flow curves. The $E(\alpha)/R$
31 value is determined from the slope of the line generated by plotting $\ln\left(\frac{d\alpha}{dt}\right)$ against $1/T(t)$
32 at a given conversion rate. Figure 5 plots the activation energy $E(\alpha)$ and $\ln A(\alpha)f(\alpha)$
33 values as functions of the reaction progress. The kinetic triplet ($f(\alpha)$, E and $\ln A$) in Eq. (3)
34 should be constant, assuming a single-step reaction. However, a number of the reactions
35 included in AN decomposition involve multiple steps that contribute to the overall reaction
36 rate measured in thermal analysis experiments. If a process involves several steps with
37 different activation energies, the overall reaction rate will vary with both temperature and
38 extent of conversion.

39 The $E(\alpha)$ and $\ln A(\alpha)f(\alpha)$ plots in Figure 4 demonstrate that both values tend to remain
40 constant with increasing α ($0.1 < \alpha < 0.5$), and $E(\alpha)$ exhibits a range of approximately
41 165-168 kJ mol⁻¹. The activation energy reported for the liquid-phase thermal decomposition
42 of AN is 170 kJ mol⁻¹ [19]. The calculated activation energy is good agreement with the
43 experimental one, and this supports the validity of the YNU-L 1.0 model.

44 This range of activation energy values does not match that of the key elementary reaction:
45 HNO₃ + HNO₃ → N₂O₅ + H₂O (R6 in Table 1, is the rate determining step based on an
46 analysis of the rate of production using the CHEMKIN-PRO software [28]). The rate of
47 production analysis also found that the major reactions that follow R6 are R7, R18 and R22,
48 and the activation energy values of these reactions do not match the apparent activation
49 energy from the Friedman method. This result is quite natural because the apparent kinetic
50 triplet ($f(\alpha)$, E and $\ln A$) is influenced by every elementary reaction composed detailed
51 kinetic model, and the model in this study is quite complicated.




Figure 1 Heat flow curves at 5 K min⁻¹ obtained from the a) P-DSC and b) YNU L1.0 simulations.




Figure 2 Variations in the AN decomposition species over temperature at a heating rate of 5 K min⁻¹.




Figure 3 Heat flow curves obtained from YNU L1.0 simulations at heating rates of 1, 2, 4 and 8 K min⁻¹.




Figure 4 Iso-conversional plots of Friedman method for heat flow curves shown in Figure 2.




Figure 5 Activation energy as a function of the reaction progress as determined based on the Friedman method for heat flow curves from YNU-L1.0 model.

Based on kinetics, we simulated the adiabatic temperature rise obtained using initial temperatures of 423, 523, 623 and 823 K. In addition, we simulated the thermal behavior at these same temperatures based on the YNU-L 1.0 model, employing the CHEMKIN-PRO software. Figure 5 shows the predicted temperature changes over time, which exhibit an abrupt rise in temperature following an induction time, clearly indicating a runaway reaction. This figure also shows time to maximum rate under adiabatic condition (TMR_{ad}) values for the various initial temperatures. The YNU-L 1.0 model predictions include higher adiabatic temperature increases than those obtained from kinetic analysis. These differences are ascribed to variations in the decomposition products. The ionic condensed-phase decomposition of AN is known to generate N_2O and H_2O . Figure 2 plots the generation of products at a heating rate of 5 K min^{-1} from 443 to 623 K (the associated heat flow profile is shown in Figure 1). Within this temperature range, the decomposition yields primarily N_2O and H_2O with only a small amount of N_2 . In contrast, at higher temperatures, AN decomposes to yield N_2 and H_2O via radical decomposition. As discussed in the Introduction, the ionic decomposition produces N_2O and H_2O . Figure 7 plots the products obtained under the adiabatic conditions at an initial temperature of 523 K. The mole number of N_2O gradually increases as the temperature rises (the associated temperature profile is shown in Figure 5a). Following the TMR_{ad} , the temperature sharply increases and N_2 gas is also evolved simultaneously. In the YNU-L 1.0 model, AN can decompose to N_2 and H_2O at high temperatures based on the radical reactions listed in Table 2. Due to the high chemical stability of N_2 , the formation of N_2 produces more heat than N_2O generation. The original exotherms elaborated in the model-free Friedman model (Figure 3) were primarily the result of the generation of N_2O rather than N_2 . Thus, the net heat of reaction and adiabatic temperature changes obtained using the model-free Friedman model are smaller than those obtained from the YNU-L 1.0 model.

Although the heat of reaction for the runaway decomposition given by the model-free Friedman model is underestimated, the model does accurately predict TMR_{ad} (within one order of magnitude of error) at initial temperatures of 423, 523 and 623 K. Exothermic thermal decomposition to N_2O and H_2O increases the temperature until the runaway reaction begins and there is an abrupt rise in temperature. Figure 7 demonstrates that N_2O and H_2O are produced during the induction time (via ionic decomposition) rather than N_2 (via radical decomposition) prior to the runaway reaction. Because the major reaction in the model-free Friedman model is the same as that during the induction reaction (representing ionic decomposition), the predictions obtained from the model-free Friedman model for the TMR_{ad} values at initial temperatures of 423, 523 and 623 K are accurate.

The difference between the TMR_{ad} values tend to gradually increase with increasing or

decreasing from decomposition onset-temperature of approximately 500 K. Especially, The difference between the TMR_{ad} values generated by the model-free Friedman model and the YNU-L1.0 model at 823 K is greater than one order of magnitude, as shown in Figure 6d). Figure 8 presents a plot of product formation under the adiabatic conditions at an initial temperature of 823 K. Here, the amounts of both N_2O and N_2 gradually increase over time as the temperature rises (the associated temperature profile is shown in Figure 6d)). More N_2 is generated compared to N_2O during the induction time, meaning that both ionic and radical decomposition take place during the TMR_{ad} span, and that the rate determining reaction in the radical mechanism is the homolytic cleavage of HNO_3 (R28 in Table 2). The primary decomposition process changes from ionic to radical with increasing temperature and this leads to the difference in the TMR_{ad} values obtained from the model-free Friedman model and YNU-L 1.0 model at 823 K. This result demonstrates that kinetic analysis does not work well if the reaction employed in the thermal analysis is different from the reaction under the associated experimental conditions.

Thermal explosion of AN is important phenomenon to consider the safe development and use of AN-containing devices. The temporal temperature profiles in Figure 6 shows a thermal explosion behavior of AN at various initial temperatures. The temperature gradually increases during the induction period, and then the temperature sharply rises (the ignition). In the induction period at initial temperature of 523 showed in Figure 7, the mole number of N_2O increases at first, and then N_2 is produced. Production of N_2O is the result of ionic mechanism discussed above, and production of N_2 is the result of radical mechanism in Table 2. The ionic reaction of AN occurs during the induction period, and AN mainly decompose to N_2O and H_2O with increasing the temperature. As a result of increasing in temperature, the radical mechanism overwhelms the ionic reaction. Then followed chain growth reactions yield N_2 , H_2O with larger heat of reaction. Although The YNU-L1.0 model successfully simulates a thermal explosion behavior of AN, the model does not include physical changes, evaporation and gasification. Further study is needed to improve the model and to obtain precise prediction on the thermal explosion of AN.




Figure 6 Temporal profiles of Temperatures simulated from i) model-free Friedman model and ii) YNU-L 1.0 model at various initial temperatures.




Figure 7 Temporal profiles of N_2 and N_2O from AN decomposition at initial temperature of 523 K. This profile was simulated based on the YNU-L 1.0 model.




Figure 8 Temporal profiles of N_2 and N_2O from AN decomposition at initial temperature of 823 K. This profile was simulated based on the YNU-L 1.0 model.

Conclusions

We analyzed the kinetics of the thermal decomposition of liquid AN based on results obtained from detailed reaction simulations. The model employed included both ionic and radical reaction mechanisms associated with AN. Rate coefficients were calculated to allow TS theory analyses of the reactions identified in a previous study. The rate coefficients for radical recombination reactions and proton transfers with no energy barriers were set to the diffusion limited value of $10^{-9} \text{ cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. Thermal correction, entropy and heat capacity values were then calculated from the partition function using statistical machinery. The heats of formation for gas-phase molecules were determined by the traditional atomization method combined

1 with the G4 level of theory. The heats of formation of molecules in solution were obtained
2 from the gas-phase heats of formation and the enthalpies of solvation at 298.15 K. An *ab*
3 *initio* model was used to simulate thermal behavior (i.e., heat flow) during AN decomposition.
4 The simulated heat flow curves were elaborated by the Friedman method. The activation
5 energy for the exothermic decomposition had a range of approximately 162-168 kJ mol⁻¹ with
6 α values between 0.1 and 0.5. The activation energy reported for the liquid-phase thermal
7 decomposition of AN is 170 kJ mol⁻¹ [19], and this agreement supports validity of the YNU-L
8 1.0 model. Calculations of TMR_{ad} based on the Friedman kinetic model provided accurate
9 predictions below 623 K. There was a difference of more than one order of magnitude
10 between the TMR_{ad} values obtained from the model-free Friedman model and YNU-L 1.0
11 model approaches at 823 K. Analyses of variations in the evolved products indicated that the
12 dominant mechanism transitions from ionic to radical decomposition with temperature
13 increase. Thermal behavior predictions based on kinetic analysis using thermal analysis are
14 evidently accurate when the dominant reaction in the thermal analysis is the same as the
15 reaction under the experimental conditions. TMR_{ad}-based studies using the model-free
16 kinetics were found to generate errors when the dominant reaction mechanism changes with
17 temperature. These errors should be taken into consideration when TMR_{ad} values are being
18 calculated to evaluate the safe operational temperatures of chemicals based on kinetic data.

19 Detailed chemical reaction simulations revealed that the thermal explosion occurs after an
20 induction period. The ionic reaction, $\text{AN} \rightarrow \text{N}_2\text{O} + 2\text{H}_2\text{O}$, starts the decomposition reaction
21 with exothermic heat during the induction period and the temperature increases gradually. As
22 a result of increasing in temperature, the radical mechanism overwhelms the ionic mechanism.
23 After radical initiation, a thermal explosion is ignited, and the temperature rises sharply.

24 Acknowledgement

25 This research was supported by JSPS KAKENHI Grant Number 17H00844.

26 Reference

- 27 1. Oxley JC, Smith JL, Wang W. Compatibility of ammonium nitrate with monomolecular
28 explosives. Part II nitroarenes. *J Phys Chem.* 1994;98:3901-7.
- 29 2. Marlair G, Kordek M. Safety and security issues relating to low capacity storage of AN-based
30 fertilizers. *J Hazard Mater.* 2005;A123:13-28.
- 31 3. U.S. chemical safety and hazard investigation board. Investigation report, final, West fertilizer
32 company fire and explosion, REPORT 2013-02-I-TX.
- 33 4. Dechy N, Bourdeaux T, Ayrault A, Kordek M, Coze JL. First lessons of the Toulouse
34 ammonium nitrate disaster, 21st September 2001, AZF plant, France. *J Hazard Mater.* 2004;
35 11:131-8.
- 36 5. Oommen C, Jain SR. Ammonium nitrate: a promising rocket propellant oxidizer. *J Hazard*
37 *Mater.* 1999;A67:253-81.
- 38 6. Chaturvedi S, Dave PN. Review on thermal decomposition of ammonium nitrate. *J Energ*
39 *Mater.* 2013;31:1-26.
- 40 7. Sinditskii VP, Egorshchikov VY, Levshenkov AI, Serushkin VV. Ammonium nitrate: combustion
41 mechanism and the role of additives. *Propellants Explos Pyrotech.* 2005;30:269-80.
- 42 8. Manelis GB, Nazin GM, Rubtsov YI, Strunin VA. Thermal decomposition and combustion of
43 explosives and propellants. CRC Press: Boca Raton; 2003.
- 44 9. Koroban VA, Burtsev YN, Alimov FR, Haustov AD, Dubovik VA, Teselkin VA, Thermal
45 decomposition features of ammonium nitrate and its boron mixture under high pressures.
46 *Propellants Explos Pyrotech.* 2004;19:307-10.
- 47 10. Cagnina S, Rotureau P, Singh S, Turcotte R, Fayet G, Adamo C. Theoretical and experimental
48 study of the reaction between ammonium nitrate and sodium salts. *Ind Eng Chem Res.*
49 2016;55:12183-90.
- 50 11. Kajiyama K, Izato Y, Miyake A. Thermal characteristics of ammonium nitrate, carbon, and
51 copper (II) oxide mixtures. *J Therm Anal Calorim.* 2013;113:1475-80.
- 52 12. Izato Y, Date S, Miyake A. Combustion characteristics of ammonium nitrate and carbon
53 mixtures based on a thermal decomposition mechanism. *Propellants Explos Pyrotech.*
54 2013;38:129-35.
- 55 13. Brower KR, Oxley JC, Tewari MP. Evidence for hemolytic decomposition of ammonium
56 nitrate at high temperature. *J Phys Chem.* 1989;93:4029-33.
- 57 14. Park J, Lin MC. Thermal decomposition of gaseous ammonium nitrate at low pressure: kinetic
58 modeling of product formation and heterogeneous decomposition of nitric acid. *J Phys Chem*
59
60
61
62
63
64
65

A. 2009;113:13556-61.

15. Skarlis SA, Nicolle A, Berthout D, Duajardin C, Granger P, Combined experimental and kinetic modeling approaches of ammonium nitrate thermal decomposition. *Thermochimica Acta*. 2014;584:58-66.
16. Izato Y, Koshi M, Miyake A. Identification of thermal decomposition products and reactions for liquid ammonium nitrate on the basis of ab initio calculation. *Int J Chem Kinet*. 2016;49:83-99.
17. Chai JD, Gordon MH. Long-range corrected hybrid density functionals with damped atom-atom dispersion corrections. *Phys Chem Chem Phys*. 2008;10:6615-20.
18. Montgomery JA, Frisch MJ, Ochterski JW, Petersson GA. A complete basis set model chemistry. VI. Use of density functional geometries and frequencies. *J Chem Phys* 1999;110:2822-7.
19. Willis A, Rosser S, Inami H, Wise H. The Kinetics of decomposition of liquid ammonium nitrate. *J Phys Chem*. 1963;67:1753-7.
20. Izato Y, Miyake A. A condensed phase decomposition mechanism for ammonium nitrate. *Sci Technol Energy Materl*. 2015;76:98-103
21. Vyazovkin S, Clawson JS, Wight CA. Thermal dissociation kinetics of solid and liquid ammonium nitrate. *Chem Mater*. 2001;13:960-6.
22. Izato Y, Miyake A. Identification of radical reactions and products for aqueous hydroxylamine nitrate (HAN) solution based on ab initio calculations. *Sci Technol Energy Materl*. 2018 in press
23. Miyoshi A. GPOP software, rev. 2013.07.15m7, available from the author. See <http://akrmys.com/gpop/>.
24. Nikolaidis A, Rauk A, Glukhovtsev MN, Radom L. Heats of formation from G2, G2(MP2), and G2(MP2,SVP) total energies. *J Phys Chem*. 1996;100:17460-564.
25. Curtiss LA, Redfern PC, Raghavachari K. Gaussian-4 theory. *J. Chem. Phys*. 2007;126:1-12.
26. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Mennucci B, Petersson GA, Nakatsuji H, Caricato M, Li X, Hratchian HP, Izmaylov AF, Bloino J, Zheng G, Sonnenberg JL, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven T, Montgomery JA, Peralta JE, Ogliaro F, Bearpark M, Heyd JJ, Brothers E, Kudin KN, Staroverov VN, Keith T, Kobayashi R, Normand J, Raghavachari K, Rendell A, Burant JC, Iyengar SS, Tomasi J, Cossi M, Rega N, Millam JM, Klene M, Knox JE, Cross JB, Bakken V, Adamo C, Jaramillo J, Gomperts R, Stratmann RE, Yazyev O, Austin AJ, Cammi R, Pomelli C, Ochterski JW, Martin RL, Morokuma K, Zakrzewski VG, Voth GA, Salvador P, Dannenberg JJ, Dapprich S, Daniels AD, Farkas O, Foresman JB, Ortiz JV, Cioslowski J, Fox DJ. *Gaussian 09, Revision C.01*. Gaussian, Inc., Wallingford CT; 2010
27. Yamashita A, Asai K. Dielectric anomaly of ammonium nitrate in low frequency region. *J Phys Soc Japan*. 1963;18:1247-53.
28. CHEMKIN-PRO 15131, Reaction Design: San Diego, 2013.
29. Friedman HL. Kinetics of thermal degradation of char-forming plastics from thermogravimetry. Application to a phenolic plastic, *J Polym Sci C*. 1963;6:183-95.
30. Ozawa T. Applicability of friedman plot. *Journal of Thermal Analysis*. 1986;31:547-51.
31. Izato Y, Miyake A. Thermal decomposition of molten ammonium nitrate (AN). *J Therm Anal Calorim* (2015) 122:595-600